



PLG gene

plasminogen

Normal Function

The *PLG* gene provides instructions for making a protein called plasminogen, which is produced in the liver. Enzymes called plasminogen activators convert plasminogen into the protein plasmin, which breaks down another protein called fibrin. Fibrin is the main protein involved in blood clots and is important for wound healing, creating the framework for normal tissue to grow back. Excess fibrin is broken down when no longer needed, and the new, more flexible normal tissue takes its place.

Health Conditions Related to Genetic Changes

congenital plasminogen deficiency

More than 50 mutations in the *PLG* gene have been identified in people with congenital plasminogen deficiency. This disorder is characterized by hard growths on the mucous membranes, which are the moist tissues that line body openings such as the eyelids and the inside of the mouth. Congenital plasminogen deficiency most often affects the conjunctiva, which are the mucous membranes that protect the white part of the eye (the sclera) and line the eyelids.

PLG gene mutations can decrease the amount of plasminogen produced, its function, or both. When the mutations affect plasminogen levels as well as the activity of the protein, affected individuals may be said to have type I congenital plasminogen deficiency, characterized by the growths previously described. People with *PLG* gene mutations that result in normal levels of plasminogen with reduced activity are said to have type II congenital plasminogen deficiency or dysplasminogenemia. This form of the condition often has no symptoms.


A reduction in functional plasminogen results in less plasmin to break down fibrin, leading to a buildup of fibrin. The excess fibrin and the resulting inflammation of the tissue result in the inflamed growths characteristic of congenital plasminogen deficiency.

It is unclear why the excess fibrin builds up in the mucous membranes but does not usually result in abnormal clots in the blood vessels (thromboses). Researchers suggest that other enzymes in the blood may also break down fibrin, helping to compensate for the reduced plasminogen levels.

Chromosomal Location

Cytogenetic Location: 6q26, which is the long (q) arm of chromosome 6 at position 26

Molecular Location: base pairs 160,702,193 to 160,754,054 on chromosome 6 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- DKFZp779M0222
- plasmin
- plasminogen isoform 1 precursor
- plasminogen isoform 2 precursor
- PLMN_HUMAN

Additional Information & Resources

Educational Resources

- Holland-Frei Cancer Medicine (sixth edition, 2003): Physiology of Normal Homeostasis
<https://www.ncbi.nlm.nih.gov/books/NBK13422/>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28PLG%5BTI%5D%29+OR+%28plasminogen%5BTI%5D%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+360+days%22%5Bdp%5D>

OMIM

- PLASMINOGEN
<http://omim.org/entry/173350>

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology
http://atlasgeneticsoncology.org/Genes/GC_PLG.html
- ClinVar
<https://www.ncbi.nlm.nih.gov/clinvar?term=PLG%5Bgene%5D>
- HGNC Gene Symbol Report
http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=9071
- NCBI Gene
<https://www.ncbi.nlm.nih.gov/gene/5340>
- UniProt
<http://www.uniprot.org/uniprot/P00747>

Sources for This Summary

- Klammt J, Kobelt L, Aktas D, Durak I, Gokbuget A, Hughes Q, Irkec M, Kurtulus I, Lapi E, Mechoulam H, Mendoza-Londono R, Palumbo JS, Steitzer H, Tabbara KF, Ozbek Z, Pucci N, Sotomayor T, Sturm M, Drogies T, Ziegler M, Schuster V. Identification of three novel plasminogen (PLG) gene mutations in a series of 23 patients with low PLG activity. *Thromb Haemost.* 2011 Mar; 105(3):454-60. doi: 10.1160/TH10-04-0216. Epub 2010 Dec 21.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/21174000>
- Mehta R, Shapiro AD. Plasminogen deficiency. *Haemophilia.* 2008 Nov;14(6):1261-8. doi: 10.1111/j.1365-2516.2008.01825.x. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/19141167>
- OMIM: PLASMINOGEN
<http://omim.org/entry/173350>
- Schuster V, Hügler B, Tefs K. Plasminogen deficiency. *J Thromb Haemost.* 2007 Dec;5(12):2315-22. Epub 2007 Sep 26. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/17900274>
- Schuster V, Seregard S. Ligneous conjunctivitis. *Surv Ophthalmol.* 2003 Jul-Aug;48(4):369-88. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/12850227>
- Tefs K, Gueorguieva M, Klammt J, Allen CM, Aktas D, Anlar FY, Aydogdu SD, Brown D, Ciftci E, Contarini P, Dempfle CE, Dostalek M, Eisert S, Gökbuget A, Günhan O, Hidayat AA, Hügler B, Isikoglu M, Irkec M, Joss SK, Klebe S, Kneppo C, Kurtulus I, Mehta RP, Ornek K, Schneppenheim R, Seregard S, Sweeney E, Turtschi S, Veres G, Zeitler P, Ziegler M, Schuster V. Molecular and clinical spectrum of type I plasminogen deficiency: A series of 50 patients. *Blood.* 2006 Nov 1; 108(9):3021-6. Epub 2006 Jul 18.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16849641>

Reprinted from Genetics Home Reference:
<https://ghr.nlm.nih.gov/gene/PLG>

Reviewed: August 2012
Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services